

Maternal Recognition of Pregnancy in Mammals

Abhay Kumar Meena^{*1}, Hitesh Kumar², Lakshmi Yadav³, Shivani Purohit⁴ And Kuldeep Singh Gurjar⁵

^{*1} Assistant professor, Livestock Research Station, Kodamdesar

^{2,3,4,5} Department of Veterinary Gynaecology and Obstetrics,

Post Graduate Institute of Veterinary Education and Research (PGIVER), Jaipur

Rajasthan University of Veterinary and Animal Sciences, Bikaner- 334001 (Rajasthan)

<https://doi.org/10.5281/zenodo.10825436>

Abstract

Maternal recognition of pregnancy refers to the requirement for the conceptus (embryo and its associated extra-embryonic membranes) to produce a hormone that acts on the uterus and/or corpus luteum (CL) to ensure maintenance of a functional CL for production of progesterone; the hormone required for pregnancy in most mammals.

1. Introduction

Maternal recognition of pregnancy (MRP) is the biological process that allows a female to recognize the presence of a conceptus and remodel the endocrine, histological, and secretory function within the reproductive tract to allow attachment/implantation and further embryonic and fetal development. This process, which can be summarized as a mechanism of prevention of luteolysis, involves signaling between the embryo and the uterus leading to successful attachment/implantation. In-depth studies on MRP have been sparked by the impact of early pregnancy loss on productivity in various species such as cattle (Diskin and Morris, 2008), camelids (Tibary and Pearson, 2015), and horses (Canisso et al., 2013; Vanderwall, 2008).

Establishment of pregnancy involves interactions between two interdependent systems defined as:

- Uterus, and
- Conceptus (embryo and extra embryonic membranes)

- At the appropriate time, the conceptus must produce steroid hormones and /or proteins to signal its presence to the maternal system.
- This signal is necessary for corpus luteum (CL) maintenance, production of progesterone and continued endometrial development and secretory activity.
- This phenomenon was described by Short (1969) as “Maternal Recognition of Pregnancy” (MRP).
- In most of the species, the conceptus provides a timely biochemical signal or pregnancy will terminate



- The critical series of events by which the Conceptus initially signals its presence to the dam and enables the pregnancy to continue is referred to as maternal recognition of pregnancy (MRP)
- MRP must occur prior to luteolysis
- If the conceptus fails to signal its presence at exactly the correct time, the function of CL is terminated by the luteolytic action of prostaglandin F₂ alpha (PGF₂ alpha) from the uterus. This ensures that the female will return to estrus and mate at frequent intervals until a successful pregnancy is established.
- Uterine PGF₂ alpha is produced by endometrium of cows, ewes, mares and sows and causes morphologic regression of CL and cessation of progesterone production.
- The effect of conceptus is luteostatic, since progesterone production is maintained at a level comparable to that of dioestrus during pregnancy.
- Basal secretion of luteinizing hormone (LH) from the anterior pituitary is also essential for CL maintenance and function during pregnancy.

2. Agents of MRP

2.1 MRP in ruminants


The MRP in ruminants is IFN τ . IFN τ acts on the endometrium in a paracrine manner to prevent luteolysis, thereby maintaining the CL and production of progesterone (Roberts, 1989b).

Originally, it was known as ovine trophoblast protein in sheep and bovine trophoblast protein in cattle, the name interferon Tau was given to it by International Interferon Society. There are different types of IFN and IFN τ is unique to ruminants. IFN τ is a Type I interferon with potential antiviral, anti-proliferative and some immunomodulatory biological activities. This implies that apart from being the signal for MRP, IFN τ plays other roles in early pregnancy in ruminants. These include protection of the conceptus or uterus against viral infection and modulating the maternal immune response towards tolerating the apparently foetal 'semi allograft'.

The transfer of trophoblastic vesicles in cyclic ewes culminates into maintenance of the CL (Martalet al. 1979). Therefore, IFN τ originates from the conceptus and not the endometrium. In sheep, secretion of IFN τ is observed from days 10 to day 21 (day 0 = mating) with maximum levels between days 16 and 18. Likely by days 22–23, the synthesis of IFN τ has ceased. The size of the blastocyst is a factor that determines the amount of IFN τ produced which may not necessarily tally with the quantity of IFN τ mRNA expressed by the blastocyst. This phenomenon was observed in cow (Robinson et al., 2006). Ovine embryos in vitro do not expand after hatching; however, they were reported to produce a small amount of IFN τ . IFN τ injection on days 11–15 post-oestrus decreased concentrations of oestrogen and progesterone receptors (PGR) in sheep endometrium on day 16 when compared with serum protein-infused control ewes, whose corpora lutea were undergoing regression on days 14–16 (Mirando et al., 1993).



Interferon from the Conceptus prevents luteolysis in the cow and Ewe



Interferon tau also causes production of proteins from the uterine glands which migrates to uterine lumen to nourish the Conceptus

2.2 MRP in sows

In sow, the MRP signal is oestrogen. For the pregnancy to be sustained by the action of oestrogen, at least two embryos, one in each of the uterine horns are required. Otherwise, the horn with no embryos will still produce $\text{PGF2}\alpha$ and this will initiate embryo loss (Senger, 2005). Pig conceptus produces oestrogen on day 14 and 18 as agent of MRP. The oestrogen redirects the $\text{PGF2}\alpha$ synthesis towards the endometrial lumen (exocrine) rather than diffusing into the blood (endocrine) (Bazer, 2013). Therefore, $\text{PGF2}\alpha$, produced by the endometrium fails to get to the ovary to cause luteolysis. In the endometrial lumen, $\text{PGF2}\alpha$ is metabolised and/or speculated to cause (local) contraction in the lumen. The latter is envisaged to aid even distribution of the embryos (being multiparous) within the available space in the uterus.

2.3 MRP in mare

In mares, the agent of MRP remains a mystery and yet to be clarified (Klein and Troedsson, 2011). Some scholars believe MRP in mare involves some glycoproteins, oestrogen and PGE2 even though these could not be substantiated (Klein and Troedsson, 2011). The study of Wilsher and Allen (2011) disproved oestrogen from being a likely agent for MRP in mares. Instead, administration of fractionated coconut or peanut oil on Day 10 post ovulation provides an effective and practical method of extending luteal life span. What is certain in this species is the need for movement of the embryo from one horn to another (McDowell et al., 1988; Sharp et al., 1989). This movement is accomplished by simple peristalsis within the endometrium and the bi-partite shape of the mare uterus seems a physiological facilitator of this movement. The simulation of embryo movement from one horn to the other using a bead or ball procrastinates



luteolysis and extends CL longevity (Rivera del alamo et al., 2008). Both conceptus and endometrium synthesised PGE₂ during early pregnancy (Boerboom et al., 2004). It was then proposed that PGE₂ by antagonising the luteolytic effect of PGF_{2a} is the MRP in this species. This speculation was further corroborated by the study of Ealy et al. (2010). The presence of conceptus on day 15 pregnancy blocked the induction of COX-2, that determines the synthesis of PGs. Sequel to this is reduced expression of PGF_{2α} and abrogation of luteolysis (Boerboom et al., 2004). This evidence remains to be clarified.

Unlike ruminant, horse conceptus or the uterine flushing on days 13–15 gestation failed to show antiviral activities of interferon. Rather, the study of Herrler et al. (2000) confirmed the insulin growth factor 1 (IGF-I) binding activity and several IGF binding proteins in the pre-implantation equine conceptus. These are hypothesised to be essentially required to enhance embryo development and so, indirectly partake in MRP.

2.4 MRP in dogs and cats

In dogs and cats, there is dearth of information on MRP or early pregnancy factor like on other species. This may be attributed to the fact that MRP is not required for sustainance of pregnancy in these species. Whether is a conceptive or nonconceptive, CL is retained for the same period of time (about 60 days). This culminates into exhibition of pseudo-pregnancy (Senger, 2005).

2.5 MRP in human

In human, the luteolytic agent is also PGF_{2a} which is produced by the ovary instead of the endometrium in ruminant and pig (Aplin et al., 2008). The signal for MRP is hCG (Ross, 1978). hCG is luteotrophic and is produced by the blastocyst on days 4–5 as soon as the embryo descends from the oviduct into the endometrium. This explains the reason why detection of hCG in the serum/urine has been used as one of the earliest pregnancy diagnosis in woman (Johnson et al., 2009).

Table 1 Time of production and agents of maternal recognition of pregnancy (MRP) in mammals.

Species	Gestation length (days)	Placental takeover (days)	Agent of MRP	Day of production
Ewe	150	50	oIFN τ	9/10–21
Cow	270	180–240	bIFN τ	12–38
Sow	114	11–30	Oestradiol	11–30
Human	270	60–70	hCG	11
Mare	330	70	Proteins/oestrogen?	14–16; 15–30
Dog, cat	60	–	None	–

References

- Aplin, J.D., Fazleabas, A.T., Glasser, S.R., Giudice, I.C., 2008. The Endometrium. Molecular, Cellular and Clinical Perspective. Informa healthcare Ltd., London, United Kingdom.
- Bazer, F.W., 2013. Pregnancy recognition signaling mechanisms in ruminants and pigs. *J. Anim. Sci. Biotechnol.* 4, 23.



- Boerboom, D., Brown, K.A., Vaillancourt, D., Poitras, P., Goff, A.K., Watanabe, K., Dore', M., Sirois, J., 2004. Expression of key prostaglandin synthases in equine endometrium during late diestrus and early pregnancy. *Biol. Reprod.* 70 (2), 391–399.
- Canisso IF, Beltaire KA, Bedford-Guaus SJ. Premature luteal regression in a pregnant mare and subsequent pregnancy maintenance with the use of oral altrenogest. *Equine Veterinary Journal* 2013; 45:97-100.
- Diskin MG, Morris DG. Embryonic and early foetal losses in cattle and other ruminants. *Reprod Domestic Anim* 2008; 43:260-267.
- Ealy, A.D., Eroh, M.L., Sharp, D.C., 2010. Prostaglandin H synthase Type 2 is differentially expressed in endometrium based on pregnancy status in pony mares and responds to oxytocin and conceptus secretions in explant culture. *Anim. Reprod. Sci.* 117, 99–105.
- Herrler, A., Pell, J.M., Allen, W.R., Beier, H.M., Stewart, F., 2000. Horse conceptuses secrete insulin-like growth factor-binding protein 3. *Biol. Reprod.* 62 (6), 1804–1811.
- Johnson, S.R., Miro, F., Barrett, S., Ellis, J.E., 2009. Levels of urinary human chorionic gonadotrophin (hCG) following conception and variability of menstrual cycle length in a cohort of women attempting to conceive. *Curr. Med. Res. Opin.* 25, 741–748.
- Klein, C., Troedsson, M.H.T., 2011. Maternal recognition of pregnancy in the horse: a mystery still to be solved. *Reprod. Fertil. Dev.* 23, 952.
- Martal, J., Lacroix, M.C., Loudes, C., Saunier, M., Wintenbergertorres, S., 1979. Trophoblastin, an antiluteolytic protein present in early pregnancy in sheep. *J. Reprod. Fertil.* 56, 63–73.
- Mcdowell, K.J., Sharp, D.C., Grubaugh, W., Thatcher, W.W., Wilcox, C.J., 1988. Restricted conceptus mobility results in failure of pregnancy maintenance in mares. *Biol. Reprod.* 39, 340–348.
- Mirando, M.A., Harney, J.P., Zhou, Y., Ogle, T.F., Ott, T.L., Moffatt, R.J., Bazer, F.W., 1993. Changes in progesterone and oestrogen receptor mRNA and protein and oxytocin receptors in endometrium of ewes after intrauterine injection of ovine trophoblast interferon. *J. Mol. Endocrinol.* 10, 185–192.
- Rivera del alamo, M.M., Reilas, T., Kindahl, H., Katila, T., 2008. Mechanisms behind intrauterine device-induced luteal persistence in mares. *Anim. Reprod. Sci.* 107, 94–106.
- Roberts, R.M., 1989b. A novel group of interferons associated with the early ovine and bovine embryo. *J. Interferon Res.* 9, 373–378.
- Robinson, R.S., Fray, M.D., Wathes, D.C., Lamming, G.E., Mann, G.E., 2006. In vivo expression of interferon tau mRNA by the embryonic trophoblast and uterine concentrations of interferon tau protein during early pregnancy in the cow. *Mol. Reprod. Dev.* 73, 470–474.
- Ross, G.T., 1978. Human chorionic gonadotropin and maternal recognition of pregnancy. *Ciba Found. Symp.* 191–208.
- Senger, P.L., 2005. *Pathways to Pregnancy and Parturition*, Current Conceptions, Pullman, WA.
- Sharp, D.C., Mcdowell, K.J., Weithenauer, J., Thatcher, W.W., 1989. The continuum of events leading to maternal recognition of pregnancy in mares. *J. Reprod. Fertil. Suppl.* 37, 101–107.
- Tibary A, Pearson LK. Pregnancy loss and abortion in camelids. *Proc of First Latin American Animal Reproduction Congress, Buenos Aires, Argentina, Mrch 11-15, 2015, pages 67-87 2015.*
- Trasorras V, Chaves MG, Neild D, Gambarotta M, and Aba M, Agüero A. Embryo transfer technique: Factors affecting the viability of the corpus luteum in llamas. *Anim Reprod Sci* 2010; 121:279-285.
- Wilsher, S., Allen, W.R., 2011. Intrauterine administration of plant oils inhibits luteolysis in the mare. *Equine Vet. J.* 43, 99–105.